

The Mesoscale Alpine Programme

newsletter



MAP

no. 13a
october 2000



<i>Note of the Editors</i>	77
<i>News from the WG-Hydrology</i>	78
<i>Contributions from the MAP Community: WG-NUM report</i>	79
<i>Extended Abstracts from the MAP Meeting 2000, Bohinjska Bistrica, Slovenia</i>	81

Editors

Dr. Georg Mayr
 Institut für Meteorologie und Geophysik
 Universität Innsbruck
 Innrain 52, A-6020 Innsbruck
 Austria

Phone +43-512-507 54 59
 Secretary +43-512-507 54 51
 Fax +43-512-507 29 24
 e-mail Georg.Mayr@uibk.ac.at

Dr. Andrea Rossa
 MeteoSwiss
 Krähbühlstrasse 58, CH-8044 Zürich
 Switzerland

Phone +41-1-256 95 39
 Secretary +41-1-256 91 11
 Fax +41-1-256 92 78
 e-mail andrea.rossa@meteoswiss.ch

Publisher

MAP newsletter
 c/o MeteoSwiss
 Krähbühlstrasse 58, CH-8044 Zürich
 Switzerland

Programme Office

Secretary

Dr. Andrea Rossa
 MeteoSwiss
 Krähbühlstrasse 58, CH-8044 Zürich
 Switzerland

Phone +41-1-256 95 39
 Secretary +41-1-256 91 11
 Fax +41-1-256 92 78
 e-mail map@meteoswiss.ch

MAP Data Centre


Internet (WWW)

<http://www.map.ethz.ch>

The MAP newsletter invites short articles on MAP-related topics. **Contributions** to the MAP newsletter should be sent to the Editor Andrea Rossa. **Please deliver your text (graphs included) in a camera-ready format** (see templates at the MAP Data Centre), and be sure that figures are suitable for black and white reproduction.

Your contribution must not exceed 2 pages!

Camera-ready format:
 16.0 cm (6.3 inch)



➔ **Deadline for contributions to the MAP newsletter No. 14:**
 February 19, 2001 (to appear in March 2001) ➔

➔ **Deadline for contributions to the MAP newsletter No. 15:**
 May 8, 2001 (to appear in June 2001) ➔

Note of the Editors

Several contributions were accidentally left out of the MAP newsletter 13. They and a version with better figure reproduction of Benoit et al.'s "First look at the MC2 performance for finescale forecasting during the MAP SOP" are included in this addendum to newsletter 13.

Georg Mayr, Andrea Rossa
Editors

News from the WG-HYDrology

Roberto Ranzi, Dept. of Civil Engineering, University of Brescia, I-25123 Brescia
ranzi@bsing.ing.unibs.it

The data delivery and the reporting phase of the MAP-SOP 'Hydrological experiments', designed in the Project P3, are now in a good shape. The summary of the conducted experiments can be found, as usually, at the WG Web site <http://www.ing.unibs.it/~dic-segr/maphyd.htm>.

Soil moisture data, helicopter-borne passive microwave radiometric measurements, water level in reservoirs in the Lago Maggiore target area, the water level and outflow of the lake, stream flow and high resolution rain gauge data in the Toce valley are already available at the MAP Data Centre.

For a preliminary description of the experiments, the following collection of papers describing the field and numerical activity has been prepared and is ready in draft form (see the Table of contents below). The final version will be available soon and at that time it will be easier to draw better defined conclusions than those preliminarily presented at the MAP meeting 2000 in Slovenia.

The role of the Working Group now is to promote the dissemination of the results of the SOP and the exchange of experiences with other interested Working Groups.

■ Report

"Hydrological aspects in the Mesoscale Alpine Programme-SOP experiment", edited by R. Ranzi and B. Bacchi, Technical Report of the University of Brescia, Department of Civil Engineering, 10, 2000.

Table of Contents

1. Ranzi, R., Hydrological aspects in the Mesoscale Alpine Programme-SOP experiment: an overview.
2. Zappa M., Matzinger N., Gurtz J., Hydrological and Meteorological Measurements at Claro (CH)- Lago Maggiore Target Area in the

MAP-SOP 1999 RIVIERA experiment including first evaluation.

3. Falappi L., S. Barontini, A. Clerici, G. Grossi, E. Savoldi, R. Ranzi, Field and laboratory soil measurements in the Toce Valley (Italy), during the MAP-SOP 1999 TOCEX experiment.
4. S. Paloscia, G. Macelloni, P. Pampaloni, E. Santi, R. Ruisi Monitoring of soil moisture and vegetation biomass in the Toce Valley (Italy) by using microwave radiometry.
5. Menziani, M., S. Pugnaghi, Soil Moisture TDR Measurements at Pallanzeno - Lago Maggiore Target Area in the MAP-SOP 1999 TOCEX experiment.
6. E. Eccel, L. Sicher, Toller, G., The field and laboratory measurements of soil hydraulic properties in the MAP-SOP 1999 TOCEX experiment.
7. Obled, Ch. and A. Djerboua, Quantitative precipitation forecasts: a real time exercise during the MAP experiment
8. Grossi, G., B. Bacchi, L. Falappi, F. Gagni, R. Ranzi, Hydrometeorological monitoring in the Toce valley in the MAP-SOP 1999 TOCEX experiment (in preparation).

First look at the MC2 performance for finescale forecasting during the MAP SOP (a WG-NUM report)

R. Benoit⁽¹⁾, H. Davies⁽²⁾, C. Schär⁽³⁾, P. Binder⁽⁴⁾, S. Chamberland⁽¹⁾, M. Desgagné⁽¹⁾, C. Girard⁽¹⁾, D. Lüthi⁽²⁾, D. Maric⁽⁵⁾, E. Müller⁽⁴⁾, P. Pellerin⁽¹⁾, J. Schmidli⁽²⁾, F. Schubiger⁽⁴⁾, M. Sprenger⁽²⁾, S. Willemse⁽⁴⁾, W. Yu⁽¹⁾, E. Zala⁽⁴⁾

⁽¹⁾ Recherche en Prévision Numérique (RPN), Environment Canada, Dorval

⁽²⁾ Institute of Atmospheric Science of ETH (LAPETH), Zürich

⁽³⁾ Geographical Institute of ETH (GIETH), Zürich

⁽⁴⁾ MeteoSwiss, Zürich

⁽⁵⁾ Swiss Center for Scientific Computing (SCSC), Manno

Introduction

The configuration of the MC2 model for MAP, featuring a 3-km window over the entire Alps, described in (Benoit et al., MAP NL 11, 1999), has been run successfully during MAP to provide special guidance to the research aircrafts, to test the explicit convection modeling, and to establish a database for mesoscale process studies over the Alps. A website was established to record the day-by-day analysis of the model forecasts by the MC2 desk crew (RPN, ETHZ and MeteoSwiss) at the MOC in Innsbruck: now at www.geo.umnw.ethz.ch/research/map_mc2/.

From Sep. 7, 1999 until Nov. 15 1999, the model was run on a continuous basis on *gottardo*, the NEC SX-4 supercomputer at the Swiss Centre for Super Computing (SCSC/CSCS); results were archived for all days and all graphical products (about 1000 images per day) are online at the MAP Data Centre. The full 27 h 3-km simulation takes 4.5 h CPU time in a dedicated mode on 10 processors at a sustained speed of approximately 10 Gflops (50% of peak theoretical performance). During the MAP field phase, this computation represents the largest computational task that was conducted in Switzerland on a daily basis.

The full suite consisted of 14-3-1 km nested runs, with the 1-km targeting (~150 km moveable domain) crucial (mostly for the GWB project, P6) aircraft missions scheduled to take place the next morning.

A first look at the results

The topography chosen for the 3-km model (Fig. 1), rather smooth, ensured un-noisy and reliable runs (not a single failure during the

period): very subtle features (e.g. detailed timings of the foehn or gapflow onset) have been simulated.




Figure 1: Topography and domain of MC2-3km

The explicit modeling of convection with 3-km instead of 2-km as initially planned, appears to have been able to catch properly only the heavy precipitation zones, while underestimating the smaller amounts, resulting in a quite large areal bias. A comprehensive evaluation of the MC2 precipitation is being carried out in collaboration with MeteoSwiss, as part of an overview paper (Benoit et al., 2001); Fig. 2a compares the mean hourly precipitation at *all* ANETZ stations (over Switzerland) with that of the MC2 corresponding points. The timing of the precipitation was well predicted, with some underestimation of the precipitation intensity.

The planned hydrologic realtime simulations (driven by the MC2 rain) really took place with the collaboration of Nick Kouwen (Waterloo, Canada). Fig. 2b shows the hydrographs for the Toce and Ticino during the SOP. There were three significant events during the MAP SOP: September 20 during IOP-02, September 26 during IOP-03 and October 21 during IOP-08. The first rainfall event that resulted in a very good flood forecast. The second rainfall was centered on the Ticino river above Bellinzona

and was not predicted by MC2 for the Ticino River. The third event was a snowfall event with precipitation amounts that would have produced significant flows if it were rain instead of snow.

It took the 1-km final MC2 adaptation to fully capture the amplitude of the intense gravity waves, such as shown on Fig. 3, for the IOP15. The model predicted approximately the right wave amplitudes (with peak values of 4.2 and 3.6 m/s, respectively). Apart from IOP 2b, no GWB false alarms were raised by the MC2.

Outlook

A complete MC2-MAP archive (1 TeraByte) has now been established at RPN that contains all of the MC2 gridded output as well as the driving boundary files: scientists interested in using parts of this MAP data are invited to enquire to the author. The adequacy of 3-km for explicit convection will be further examined, as well as the question of the proper terrain definition.

A second MC2 suite, at 14 km, with 75 levels and a higher lid at 35 km, out to T+72h, driven by the ECMWF model, was also run on a DLR SX-4 machine and is not described here.

References

Benoit, R., M. Desgagné, P. Pellerin, S. Pellerin, S. Desjardins and Y. Chartier, 1997: The Canadian MC2: a semi-Lagrangian, semi-implicit wide-band atmospheric model suited

for fine-scale process studies and simulation. *Mon. Wea. Rev.*, **125**, 2382-2415.

Benoit, R. and co-authors, 1999: Realtime ultrafinescale modelling effort for the MAP field phase. MAP Newsletter no. 11, 14-15. Available at www.map.ethz.ch/newsletter.htm.

Benoit, R. and coauthors, 2001: The real-time ultrafinescale forecast support during the Special Observing Period of the MAP. BAMS article in preparation.




Figure 3. Vertical velocity observed by Electra (in solid line) at altitude of 6400 m ASL, and simulated by MC2-1km (dashed line) at an altitude of 6500 m ASL for IOP 15. Abscissa is latitude of flight track and model points.




Figure 2, (a): Averaged hourly precipitation observed by the ANETZ stations and predicted by MC2-3km (shifted down by 1 mm/h on plot). (b): Observed and modelled streamflow for the MAP-SOP for the Toce and Ticino (shifted down by 1000 m³/s) Rivers. One model driven by MC2 forecasts and the other, driven by radar observations (extracted from the Alpine radar composite imagery). Hydrometric observations shown at higher density during the wet episode of IOP 2-3 (17-29 September 1999).

Extended Abstracts from the MAP Meeting 2000, Bohinjska Bistrica, Slovenia

■ SESSION P5 Unstationary aspects of Foehn in a large valley

Lothon, M., B. Bénech, B. Campistron, and E. Richard: *The Foehn event of October 20/21 1999 observed during MAP IOP8* 82

■ SESSION P7 Potential vorticity banners

Bénech, B., H. Berger, J.-L. Attié, M. Lothon, B. Dartiguelongues, B. Terliuc, and A. Ezcurra: *First analysis of the constant volume balloon data during the intensive observing periods of MAP* 84

Bernard, S., A. Druilhet, F. Saïd, J.-L. Attié, and B. Bénech: *Contribution to airborne phase of dry-MAP during intensive observing periods* 86